

Solutions of some nonlinear parabolic equations with initial blow-up

Waad Al Sayed Laurent Véron

Laboratoire de Mathématiques et Physique Théorique,
Université François Rabelais, Tours, FRANCE

Abstract We study the existence and uniqueness of solutions of $\partial_t u - \Delta u + u^q = 0$ ($q > 1$) in $\Omega \times (0, \infty)$ where $\Omega \subset \mathbb{R}^N$ is a domain with a compact boundary, subject to the conditions $u = f \geq 0$ on $\partial\Omega \times (0, \infty)$ and the initial condition $\lim_{t \rightarrow 0} u(x, t) = \infty$. By means of Brezis' theory of maximal monotone operators in Hilbert spaces, we construct a minimal solution when $f = 0$, whatever is the regularity of the boundary of the domain. When $\partial\Omega$ satisfies the parabolic Wiener criterion and f is continuous, we construct a maximal solution and prove that it is the unique solution which blows-up at $t = 0$.

1991 Mathematics Subject Classification. 35K60.

Key words. Parabolic equations, singular solutions, semi-groups of contractions, maximal monotone operators, Wiener criterion.

1 Introduction

Let Ω be a domain of $\mathbb{R}^N$ ($N \geq 1$) with a compact boundary, $Q_\infty^\Omega = \Omega \times (0, \infty)$ and $q > 1$. This article deals with the question of the solvability of the following Cauchy-Dirichlet problem $\mathcal{P}^{\Omega, f}$

$$\begin{cases} \partial_t u - \Delta u + |u|^{q-1}u = 0 & \text{in } Q_\infty^\Omega \\ u = f & \text{on } \partial\Omega \times (0, \infty) \\ \lim_{t \rightarrow 0} u(x, t) = \infty & \forall x \in \Omega. \end{cases} \quad (1.1)$$

If no assumption of regularity is made on $\partial\Omega$, the boundary data $u = f$ cannot be prescribed in sense of continuous functions. However, the case $f = 0$ can be treated if the vanishing condition on $\partial\Omega \times (0, \infty)$ is understood in the H_0^1 local sense. We construct a positive solution $\underline{u}_\Omega$ of (1.1) with $f = 0$ belonging to $C(0, \infty; H_0^1(\Omega) \cap L^{q+1}(\Omega))$ thanks to Brezis results of contractions semigroups generated by subdifferential of proper convex functions in Hilbert spaces. We can also consider an internal increasing approximation of Ω by smooth bounded domains Ω^n such that $\Omega = \cup_n \Omega^n$. For each of these domains, there exists a maximal solution $\overline{u}_{\Omega^n}$ of problem $\mathcal{P}^{\Omega^n, 0}$. Furthermore the sequence $\{\overline{u}_{\Omega^n}\}$ is increasing. The limit function $u_\Omega := \lim_{n \rightarrow \infty} \overline{u}_{\Omega^n}$ is the natural candidate to be the minimal positive solution of a solution of $\mathcal{P}^{\Omega, 0}$. We prove that $\underline{u}_\Omega = u_\Omega$. If $\partial\Omega$ satisfies the parabolic Wiener

criterion [9], there truly exist solutions of $\mathcal{P}^{\Omega,0}$. We construct a maximal solution $\overline{u}_\Omega$ of this problem. Our main result is the following:

Theorem 1. *If $\partial\Omega$ is compact and satisfies the parabolic Wiener criterion, there holds*

$$\overline{u}_\Omega = \underline{u}_\Omega.$$

In the last section, we consider the full problem $\mathcal{P}^{\Omega,f}$. Under the same regularity and boundedness assumption on $\partial\Omega$ we construct a maximal solution $\overline{u}_{\Omega,f}$ and we prove

Theorem 2. *If $\partial\Omega$ is compact and satisfies the parabolic Wiener criterion, and if $f \in C(0, \infty; \partial\Omega)$ is nonnegative, $\overline{u}_{\Omega,f}$ is the only positive solution to problem $\mathcal{P}^{\Omega,f}$.*

These type of results are to be compared with the ones obtained by the same authors [1] in which paper the following problem is considered

$$\begin{cases} \partial_t u - \Delta u + |u|^{q-1}u = 0 & \text{in } Q_\infty^\Omega \\ \lim_{\text{dist}(x, \partial\Omega) \rightarrow 0} u(x, t) = \infty & \text{locally uniformly on } (0, \infty) \\ u(x, 0) = f & \forall x \in \Omega. \end{cases} \quad (1.2)$$

In the above mentioned paper, it is proved two types of existence and uniqueness result with $f \in L_{loc}^1(\Omega)$, $f \geq 0$: either if $\partial\Omega = \partial\overline{\Omega}^c$ and $1 < q < N/(N-2)$, or if $\partial\Omega$ is locally the graph of a continuous function and $q > 1$.

Our paper is organized as follows: 1- Introduction. 2- Minimal and maximal solutions. 3- Uniqueness of large solutions. 4- Bibliography.

2 Minimal and maximal solutions

Let $q > 1$ and Ω be a proper domain of $\mathbb{R}^N$, $N > 1$ with a non-empty compact boundary. We set $Q_\infty^\Omega = \Omega \times (0, \infty)$ and consider the following problem

$$\begin{cases} \partial_t u - \Delta u + u^q = 0 & \text{in } \Omega \times (0, \infty) \\ u(x, t) = 0 & \text{on } \partial\Omega \times (0, \infty). \end{cases} \quad (2.1)$$

If there is no regularity assumption on $\partial\Omega$, a natural way to consider the boundary condition is to impose $u(\cdot, t) \in H_0^1(\Omega)$. The Hilbertian framework for this equation has been studied by Brezis in a key article [2] (see also the monography [3] for a full treatment of related questions) in considering the maximal monotone operator $v \mapsto A(v) := -\Delta v + |v|^{q-1}v$ seen as the subdifferential of the proper lower semi-continuous function

$$J_\Omega(v) = \begin{cases} \int_\Omega \left(\frac{1}{2} |\nabla v|^2 + \frac{1}{q+1} |v|^{q+1} \right) dx & \text{if } v \in H_0^1(\Omega) \cap L^{q+1}(\Omega) \\ \infty & \text{if } v \notin H_0^1(\Omega) \cap L^{q+1}(\Omega). \end{cases} \quad (2.2)$$

In that case, the domain of $A = \partial J_\Omega$ is $D(A) := \{u \in H_0^1(\Omega) \cap L^{q+1}(\Omega) : \Delta u \in L^2(\Omega)\}$, and we endow $D_\Omega(-\Delta, \cdot)$ with the graph norm of the Laplacian in $H_0^1(\Omega)$

$$\|v\|_{D_\Omega(-\Delta)} = \left(\int_\Omega ((\Delta v)^2 + |\nabla v|^2 + v^2) dx \right)^{1/2}.$$

Brezis' result is the following.

Theorem 2.1 *Given $u_0 \in L^2(\Omega)$ there exists a unique function $v \in L^2_{loc}(0, \infty; D_\Omega(-\Delta)) \cap C(0, \infty; H_0^1(\Omega) \cap L^{q+1}(\Omega))$ such that $\partial_t v \in L^2_{loc}(0, \infty; L^2(\Omega))$ satisfying*

$$\begin{cases} \partial_t v - \Delta v + |v|^{q-1}v = 0 & \text{a.e. in } Q_\infty^\Omega \\ v(., 0) = u_0 & \text{a.e. in } \Omega. \end{cases} \quad (2.3)$$

Furthermore the mapping $(t, u_0) \mapsto v(t, .)$ defines an order preserving contraction semigroup in $L^2(\Omega)$, denoted by $S^{\partial J_\Omega}(t)[u_0]$, and the following estimate holds

$$\|\partial_t v(t, .)\|_{L^2(\Omega)} \leq \frac{1}{t\sqrt{2}} \|u_0\|_{L^2(\Omega)}. \quad (2.4)$$

From this result, we have only to consider solutions of (2.1) with the above regularity.

Definition 2.2 *We denote by $\mathcal{I}(Q_\infty^\Omega)$ the set of positive functions $u \in L^2_{loc}(0, \infty; D_\Omega(-\Delta)) \cap C(0, \infty; H_0^1(\Omega) \cap L^{q+1}(\Omega))$ such that $\partial_t u \in L^2_{loc}(0, \infty; L^2(\Omega))$ satisfying*

$$\partial_t u - \Delta u + |u|^{q-1}u = 0 \quad (2.5)$$

in the semigroup sense, i. e.

$$\frac{du}{dt} + \partial J_\Omega(u) = 0 \quad \text{a.e. in } (0, \infty). \quad (2.6)$$

If Ω is not bounded it is usefull to introduce another class which takes into account the Dirichlet condition on $\partial\Omega$: we assume that $\Omega^c \subset B_{R_0}$, denote by $\Omega_R = \Omega \cap B_R$ ($R \geq R_0$) and by $\tilde{H}_0^1(\Omega_R)$ the closure in $H_0^1(\Omega_R)$ of the restrictions to Ω_R of functions in $C_0^\infty(\Omega)$, thus we endow $D_{\Omega_R}(-\Delta,)$ with the graph norm of the Laplacian in $\tilde{H}_0^1(\Omega_R)$

$$\|v\|_{D_{\Omega_R}(-\Delta)} = \left(\int_{\Omega_R} ((\Delta v)^2 + |\nabla v|^2 + v^2) dx \right)^{1/2}.$$

Definition 2.3 *If Ω is not bounded but $\Omega^c \subset B_{R_0}$, we denote by $\mathcal{I}(Q_\infty^{\Omega_{loc}})$ the set of positive functions $u \in L^2_{loc}(Q_\infty^\Omega)$ such that, for any $R > R_0$, $u \in L^2_{loc}(0, \infty; D_{\Omega_R}(-\Delta)) \cap C(0, \infty; \tilde{H}_0^1(\Omega_R) \cap L^{q+1}(\Omega_R))$, $\partial_t u \in L^2_{loc}(0, \infty; L^2(\Omega_R))$ and u satisfies (2.5) in a. e. in Q_∞^Ω .*

Lemma 2.4 *If $u \in \mathcal{I}(Q_\infty^\Omega)$ or $\mathcal{I}(Q_\infty^{\Omega_{loc}})$, its extension $\tilde{u}$ by zero outside Ω is a subsolution of (2.1) in $(0, \infty) \times \mathbb{R}^N$ such that $\tilde{u} \in C(0, \infty; H_0^1(\mathbb{R}^N) \cap L^{q+1}(\mathbb{R}^N))$ and $\partial_t \tilde{u} \in L^2_{loc}(0, \infty; L^2(\mathbb{R}^N))$.*

Proof. The proof being similar in the two cases, we assume Ω bounded. We first notice that $\tilde{u} \in C(0, \infty; H_0^1(\mathbb{R}^N))$ since $\|\tilde{u}\|_{H_0^1(\mathbb{R}^N)} = \|u\|_{H_0^1(\Omega)}$. For $\delta > 0$ we set

$$P_\delta(r) = \begin{cases} r - 3\delta/2 & \text{if } r \geq 2\delta \\ r^2/2\delta - r + \delta/2 & \text{if } \delta < r < 2\delta \\ 0 & \text{if } r \leq \delta \end{cases}$$

and denote by u_δ the extension of $P_\delta(u)$ by zero outside Q_∞^Ω . Since $u_{\delta t} = P'_\delta(u)\partial_t u$, then $u_{\delta t} \in L^2_{loc}(0, \infty; L^2(\mathbb{R}^N))$ and $\|u_{\delta t}\|_{L^2} \leq \|\partial_t u\|_{L^2}$. In the same way $\nabla u_\delta = P'_\delta(u)\nabla u$, thus $u_\delta \in L^2_{loc}(0, \infty; H_0^1(\mathbb{R}^N))$ and $\|u_\delta\|_{H_0^1} \leq \|u\|_{H_0^1}$. Finally $-\Delta u_\delta = -P'_\delta(u)\Delta u - P''_\delta(u)|\nabla u|^2$. Using the fact that $P'_\delta u^q \geq u_\delta^q$, we derive from (2.6)

$$\partial_t u_\delta - \Delta u_\delta + u_\delta^q \leq 0$$

in the sense that

$$\iint_{Q_\infty^{\mathbb{R}^N}} (\partial_t u_\delta \zeta + \nabla u_\delta \cdot \nabla \zeta + u_\delta^q \zeta) dx dt \leq 0 \quad (2.7)$$

for all $\zeta \in C^\infty((0, \infty) \times \mathbb{R}^N)$, $\zeta \geq 0$. Actually, $C^\infty((0, \infty) \times \mathbb{R}^N)$ can be replaced by $L^2(\epsilon, \infty; H_0^1(\mathbb{R}^N)) \cap L^{q'}((\epsilon, \infty) \times \mathbb{R}^N)$. Letting $\delta \rightarrow 0$ and using Fatou's theorem implies that (2.7) holds with u_δ replaced by $\tilde{u}$. $\square$

Lemma 2.5 *For any $u \in \mathcal{I}(Q_\infty^\Omega)$, there holds*

$$u(x, t) \leq \left(\frac{1}{(q-1)t} \right)^{1/(q-1)} := \phi_q(t) \quad \forall (x, t) \in Q_\infty^\Omega. \quad (2.8)$$

Proof. Let $\tau > 0$. Since the function $\phi_{q,\tau}$ defined by $\phi_{q,\tau}(t) = \phi_q(t - \tau)$ is a solution of

$$\phi'_{q,\tau} + \phi_{q,\tau}^q = 0$$

and $(u - \phi_{q,\tau})_+ \in C(0, \infty; H_0^1(\Omega))$, there holds

$$\frac{1}{2} \frac{d}{dt} \int_\Omega (u - \phi_{q,\tau})_+^2 dx + \iint_{Q_\infty^\Omega} (\nabla u \cdot \nabla (u - \phi_{q,\tau})_+ + (u^q - \phi_{q,\tau}^q)(u - \phi_{q,\tau})_+) dx dt = 0.$$

Thus $s \mapsto \|(u - \phi_{q,\tau})_+(s)\|_{L^2}$ is nonincreasing. By Lebesgue's theorem,

$$\lim_{s \downarrow \tau} \|(u - \phi_{q,\tau})_+(s)\|_{L^2} = 0,$$

thus $u(x, t) \leq \phi_{q,\tau}(t)$ a.e. in Ω . Letting $\tau \downarrow 0$ and using the continuity yields to (2.8). $\square$

Theorem 2.6 *For any $q > 1$, the set $\mathcal{I}(Q_\infty^\Omega)$ admits a least upper bound $\underline{u}_\Omega$ for the order relation. If Ω is bounded, $\underline{u}_\Omega \in \mathcal{I}(Q_\infty^\Omega)$; if it is not the case, then $\underline{u}_\Omega \in \mathcal{I}(Q_\infty^{\Omega_{loc}})$.*

Proof. *Step 1- Construction of $\underline{u}_\Omega$ when Ω is bounded.* For $k \in \mathbb{N}^*$ we consider the solution $v = v_k$ (in the sense of Theorem 2.1 with the corresponding maximal operator in $L^2(\Omega)$) of

$$\begin{cases} \partial_t v - \Delta v + v^q = 0 & \text{in } \Omega \times (0, \infty) \\ v(x, t) = 0 & \text{in } \partial\Omega \times (0, \infty) \\ v(x, 0) = k & \text{in } \Omega. \end{cases} \quad (2.9)$$

When $k \rightarrow \infty$, v_k increases and converges to some $\underline{u}_\Omega$. Because of (2.8) and the fact that Ω is bounded, $\underline{u}_\Omega(t, \cdot) \in L^2(\Omega)$ for $t > 0$. It follows from the closedness of maximal

monotone operators that $\underline{u}_\Omega \in L^2_{loc}(0, \infty; D_\Omega(-\Delta)) \cap C(0, \infty; H^1_0(\Omega) \cap L^{q+1}(\Omega))$, $\partial_t \underline{u}_\Omega \in L^2_{loc}(0, \infty; L^2(\Omega))$ and

$$\frac{d\underline{u}_\Omega}{dt} + \partial J_\Omega(\underline{u}_\Omega) = 0 \quad \text{a.e. in } (0, \infty). \quad (2.10)$$

Thus $\underline{u}_\Omega \in \mathcal{I}(Q^\Omega_\infty)$. For $\tau, \epsilon > 0$, the function $t \mapsto \underline{u}_\Omega(x, t - \tau) + \epsilon$ is a supersolution of (2.1). Let $u \in \mathcal{I}(Q^\Omega_\infty)$; for $k > \phi_q(\tau)$, the function $(x, t) \mapsto (u(x, t) - \underline{u}_\Omega(x, t - \tau) - \epsilon)_+$ is a subsolution of (2.1) and belongs to $C(\tau, \infty; H^1_0(\Omega))$. Since it vanishes at $t = \tau$, it follows from Brezis' result that it is identically zero, thus $u(x, t) \leq \underline{u}_\Omega(x, t - \tau) + \epsilon$. Letting $\epsilon, \tau \downarrow 0$ implies the claim.

Step 2- Construction of $\underline{u}_\Omega$ when Ω is unbounded. We assume that $\partial\Omega \subset B_{R_0}$ and for $n > R_0$, we recall that $\Omega_n = \Omega \cap B_n$. For $k > 0$, we denote by $\underline{u}_{\Omega_n}$ the solution obtained in Step 1. Then $\underline{u}_{\Omega_n} = \lim_{k \rightarrow \infty} v_{n,k}$ where $v_{n,k}$ is the solution, in the sense of maximal operators in Ω_n of

$$\begin{cases} \frac{dv_{n,k}}{dt} + \partial J_{\Omega_n}(v_{n,k}) = 0 & \text{a.e. in } (0, \infty) \\ v_{n,k}(0) = k. \end{cases} \quad (2.11)$$

It follows from Lemma 2.4 that the extension $\tilde{v}_{n,k}$ by 0 of $v_{n,k}$ in Ω_{n+1} is a subsolution for the equation satisfied by $v_{n+1,k}$, with a smaller initial data, therefore $\tilde{v}_{n,k} \leq v_{n+1,k}$. This implies $\tilde{u}_{\Omega_n} \leq u_{\Omega_{n+1}}$. Thus we define $\underline{u}_\Omega = \lim_{n \rightarrow \infty} \underline{u}_{\Omega_n}$. It follows from Lemma 2.5 and standard regularity results for parabolic equations that $u = \underline{u}_\Omega$ satisfies

$$\partial_t u - \Delta u + u^q = 0 \quad (2.12)$$

in Q^Ω_∞ . Multiplying

$$\frac{d\underline{u}_{\Omega_n}}{dt} + \partial J_{\Omega_n}(\underline{u}_{\Omega_n}) = 0 \quad (2.13)$$

by $\eta^2 \underline{u}_{\Omega_n}$ where $\eta \in C^\infty_0(\mathbb{R}^N)$ and integrating over Ω_n , yields to

$$2^{-1} \frac{d}{dt} \int_{\Omega_n} \eta^2 \underline{u}_{\Omega_n}^2 dx + \int_{\Omega_n} \left(|\nabla \underline{u}_{\Omega_n}|^2 + \underline{u}_{\Omega_n}^{q+1} \right) \eta^2 dx + 2 \int_{\Omega_n} \nabla \underline{u}_{\Omega_n} \cdot \nabla \eta \eta \underline{u}_{\Omega_n} dx = 0.$$

Thus, by Young's inequality,

$$2^{-1} \frac{d}{dt} \int_{\Omega_n} \eta^2 \underline{u}_{\Omega_n}^2 dx + \int_{\Omega_n} \left(2^{-1} |\nabla \underline{u}_{\Omega_n}|^2 + \underline{u}_{\Omega_n}^{q+1} \right) \eta^2 dx \leq 2 \int_{\Omega_n} |\nabla \eta|^2 \underline{u}_{\Omega_n}^2 dx.$$

If we assume that $0 \leq \eta \leq 1$, $\eta = 1$ on B_R ($R > R_0$) and $\eta = 0$ on B_{2R}^c , we derive, for any $0 < \tau < t$,

$$\begin{aligned} 2^{-1} \int_{\Omega_n} \underline{u}_{\Omega_n}^2(., t) \eta^2 dx + \int_\tau^t \int_{\Omega_n} \left(2^{-1} |\nabla \underline{u}_{\Omega_n}|^2 + \underline{u}_{\Omega_n}^{q+1} \right) \eta^2 dx ds \\ \leq 2 \int_\tau^t \int_{\Omega_n} \underline{u}_{\Omega_n}^2 |\nabla \eta|^2 dx ds + 2^{-1} \int_{\Omega_n} \underline{u}_{\Omega_n}^2(., \tau) \eta^2 dx. \end{aligned} \quad (2.14)$$

From this follows, if $n > 2R$,

$$2^{-1} \int_{\Omega \cap B_R} \underline{u}_{\Omega_n}^2(., t) dx + \int_\tau^t \int_{\Omega \cap B_R} \left(2^{-1} |\nabla \underline{u}_{\Omega_n}|^2 + \underline{u}_{\Omega_n}^{q+1} \right) dx ds \leq CR^N (t+1) \tau^{-2/(q-1)}. \quad (2.15)$$

If we let $n \rightarrow \infty$ we derive by Fatou's lemma

$$2^{-1} \int_{\Omega \cap B_R} \underline{u}_\Omega^2(., t) dx + \int_\tau^t \int_{\Omega \cap B_R} \left(2^{-1} |\nabla \underline{u}_\Omega|^2 + \underline{u}_\Omega^{q+1} \right) dx ds \leq CR^N (t+1) \tau^{-2/(q-1)}. \quad (2.16)$$

For $\tau > 0$ fixed, we multiply (2.13) by $(t-\tau)\eta^2 d\underline{u}_{\Omega_n}/dt$, integrate on $(\tau, t) \times \Omega_n$ and get

$$\begin{aligned} (t-\tau) \int_{\Omega_n} \left| \frac{d\underline{u}_{\Omega_n}}{dt} \right|^2 \eta^2 dx + \frac{d}{dt} (t-\tau) \int_{\Omega_n} \left(\frac{|\nabla \underline{u}_{\Omega_n}|^2}{2} + \frac{\underline{u}_{\Omega_n}^{q+1}}{q+1} \right) \eta^2 dx \\ = \int_{\Omega_n} \left(\frac{|\nabla \underline{u}_{\Omega_n}|^2}{2} + \frac{\underline{u}_{\Omega_n}^{q+1}}{q+1} \right) \eta^2 dx - 2(t-\tau) \int_{\Omega_n} \nabla \underline{u}_{\Omega_n} \cdot \nabla \eta \frac{d\underline{u}_{\Omega_n}}{dt} \eta dx. \end{aligned}$$

Since

$$2(t-\tau) \left| \int_{\Omega_n} \nabla \underline{u}_{\Omega_n} \cdot \nabla \eta \frac{d\underline{u}_{\Omega_n}}{dt} \eta dx \right| \leq \frac{(t-\tau)}{2} \int_{\Omega_n} \left| \frac{d\underline{u}_{\Omega_n}}{dt} \right|^2 \eta^2 dx + 4(t-\tau) \int_{\Omega_n} |\nabla \underline{u}_{\Omega_n}|^2 |\nabla \eta|^2 dx,$$

we get, in assuming again $n > 2R$,

$$\begin{aligned} 2^{-1} \int_\tau^t \int_{\Omega} (s-\tau) \left| \frac{d\underline{u}_{\Omega_n}}{dt} \right|^2 \eta^2 dx ds + (t-\tau) \int_{\Omega} \left(\frac{|\nabla \underline{u}_{\Omega_n}|^2}{2} + \frac{\underline{u}_{\Omega_n}^{q+1}}{q+1} \right) \eta^2 dx \\ \leq 4 \int_\tau^t (s-\tau) \int_{\Omega} |\nabla \underline{u}_{\Omega_n}|^2 |\nabla \eta|^2 dx ds, \end{aligned} \quad (2.17)$$

from which follows,

$$\begin{aligned} 2^{-1} \int_\tau^t \int_{\Omega \cap B_R} (s-\tau) \left| \frac{d\underline{u}_{\Omega_n}}{dt} \right|^2 dx ds + (t-\tau) \int_{\Omega \cap B_R} \left(\frac{|\nabla \underline{u}_{\Omega_n}|^2}{2} + \frac{\underline{u}_{\Omega_n}^{q+1}}{q+1} \right) dx \\ \leq 4 \int_\tau^t (s-\tau) \int_{\Omega \cap B_{2R}} |\nabla \underline{u}_{\Omega_n}|^2 dx ds. \end{aligned} \quad (2.18)$$

The right-hand side of (2.18) remains uniformly bounded by $8C(2R)^N(t-\tau)t\tau^{-2/(q-1)}$ from (2.15). Then

$$\begin{aligned} 2^{-1} \int_\tau^t \int_{\Omega \cap B_R} (s-\tau) \left| \frac{d\underline{u}_{\Omega_n}}{dt} \right|^2 dx ds + (t-\tau) \int_{\Omega \cap B_R} \left(\frac{|\nabla \underline{u}_{\Omega_n}|^2}{2} + \frac{\underline{u}_{\Omega_n}^{q+1}}{q+1} \right) dx \\ \leq 8C(2R)^N(t-\tau)t\tau^{-2/(q-1)} \end{aligned} \quad (2.19)$$

By Fatou's lemma the same estimate holds if $\underline{u}_{\Omega_n}$ is replaced by $\underline{u}_\Omega$. Notice also that this estimate implies that $\underline{u}_\Omega$ vanishes in the H_0^1 -sense on $\partial\Omega$ since $\eta \underline{u}_\Omega \in H_0^1(\Omega)$ where the function $\eta \in C_0^\infty(\mathbb{R}^N)$ has value 1 in B_R and $\Omega^c \subset B_R$. Moreover estimates (2.16) and (2.19) imply that $\underline{u}_\Omega$ satisfies (2.12) a.e., and thus it belongs to $\mathcal{I}(Q_\infty^{\Omega_{loc}})$.

Step 3- Comparison. At end, assume $u \in \mathcal{I}(Q_\infty^\Omega)$. For $R > n_0$ let W_R be the maximal solution of

$$-\Delta W_R + W_R^q = 0 \quad \text{in } B_R. \quad (2.20)$$

Existence follows from Keller-Osserman's construction [5],[8], and the following scaling and blow-up estimates holds

$$W_R(x) = R^{-2/(q-1)}W_1(x/R), \quad (2.21)$$

and

$$W_R(x) = C_q(R - |x|)^{-2/(q-1)}(1 + o(1)) \text{ as } |x| \rightarrow R. \quad (2.22)$$

For $\tau > 0$ set $v(x, t) = u(x, t) - \underline{u}_\Omega(x, t - \tau) - W_R(x)$. Then v_+ is a subsolution. Since $v(\cdot, \tau) \in L^2(\Omega)$, $\lim_{s \downarrow \tau} \|v_+(\cdot, s)\|_{L^2} = 0$. Because $\eta \underline{u}_\Omega \in H_0^1(\Omega)$ for η as above, $\eta v_+ \in H_0^1(\Omega)$. Next, $\text{supp } v_+ \subset \Omega \cap B_R$. Since $u, \underline{u}_\Omega$ are locally in H^1 , we can always assume that their restrictions to $\partial B_R \times [0, T]$ are integrable for the corresponding Hausdorff measure. Therefore Green's formula is valid, which implies

$$-\int_\tau^t \int_{\Omega \cap B_R} \Delta v_+ dx dt = \int_\tau^t \int_{\Omega \cap B_R} |\nabla v_+|^2 dx dt \quad \forall t > \tau.$$

Therefore

$$\int_{\Omega \cap B_R} v_+^2(x, t) dx + \int_\tau^t \int_{\Omega \cap B_R} (|\nabla v_+|^2 + (u - (\underline{u}_\Omega(\cdot, t - \tau) + W_R)^q) v_+) dx dt \leq \int_{\Omega \cap B_R} v_+^2(x, s) dx.$$

We let $s \downarrow \tau$ and get $v_+ = 0$, equivalently $u(x, t) \leq \underline{u}_\Omega(x, t - \tau) + W_R(x)$. Then we let $R \rightarrow \infty$ and $\tau \rightarrow 0$ and obtain $u(x, t) \leq \underline{u}_\Omega(x, t)$, which is the claim. $\square$

Corollary 2.7 Assume $\Omega^1 \subset \Omega^2 \subset \mathbb{R}^N$ are open domains, then $\underline{u}_{\Omega^1} \leq \underline{u}_{\Omega^2}$. Furthermore, if $\Omega = \cup \Omega^n$ where $\Omega^n \subset \Omega^{n+1}$, then

$$\lim_{n \rightarrow \infty} \underline{u}_{\Omega^n} = \underline{u}_\Omega, \quad (2.23)$$

locally uniformly in Q_∞^Ω .

Proof. The first assertion follows from the proof of Theorem 2.6. It implies

$$\lim_{n \rightarrow \infty} \underline{u}_{\Omega^n} = u_\Omega^* \leq \underline{u}_\Omega,$$

and u_Ω^* is a positive solution of (2.5) in Q_∞^Ω . There exists a sequence $\{u_{0,m}\} \subset L^2(\Omega)$ such that $S^{\partial J_\Omega}(t)[u_{0,m}] \uparrow \underline{u}_\Omega$ as $n \rightarrow \infty$, locally uniformly in Q_∞^Ω . Set $u_{0,m,n} = u_{0,m} \chi_{\Omega^n}$; since $u_{0,m,n} \rightarrow u_{0,m}$ in $L^2(\Omega)$ then $S^{\partial J_\Omega}(\cdot)[u_{0,m,n}] \uparrow S^{\partial J_\Omega}(\cdot)[u_{0,m}]$ in $L^\infty(0, \infty; L^2(\Omega))$. If $\tilde{v}_{m,n}$ is the extension of $v_{m,n} := S^{\partial J_{\Omega^n}}(\cdot)[u_{0,m,n}]$ by zero outside $Q_\infty^{\Omega^n}$ it is a subsolution smaller than $S^{\partial J_\Omega}(\cdot)[u_{0,m,n}]$ and $n \mapsto \tilde{v}_{m,n}$ is increasing; we denote by $\tilde{v}_m$ its limit as $n \rightarrow \infty$. Since for any $\zeta \in C_0^{2,1}([0, \infty) \times \Omega)$ we have, for n large enough and $s > 0$,

$$-\int_0^s \int_\Omega (\tilde{v}_{m,n} (\partial_t \zeta + \Delta \zeta)) dx dt = \int_\Omega u_{0,m,n} \zeta(x, 0) dx - \int_\Omega \tilde{v}_{m,n}(x, s) \zeta(x, t) dx,$$

it follows

$$-\int_0^s \int_\Omega (\tilde{v}_m (\partial_t \zeta + \Delta \zeta)) dx dt = \int_\Omega u_{0,m} \zeta(x, 0) dx - \int_\Omega \tilde{v}_m(x, s) \zeta(x, t) dx.$$

Clearly $\tilde{v}_m$ is a solution of (2.5) in $Q_\infty^{\Omega^n}$. Furthermore

$$\lim_{t \rightarrow 0} \tilde{v}_m(t, \cdot) = u_{0,m} \quad \text{a.e. in } \Omega.$$

Because

$$\|\tilde{v}_m(t, \cdot) - u_{0,m}\|_{L^2(\Omega)} \leq 2 \|u_{0,m}\|_{L^2(\Omega)},$$

it follows from Lebesgue's theorem that $t \mapsto \tilde{v}_m(t, \cdot)$ is continuous in $L^2(\Omega)$ at $t = 0$. Furthermore, for any $t > 0$ and $h \in (-t, t)$, we have from 2.4 ,

$$\begin{aligned} \|\tilde{v}_{m,n}(t+h, \cdot) - \tilde{v}_{m,n}(t, \cdot)\|_{L^2(\Omega^n)} &\leq \frac{|h|}{t\sqrt{2}} \|u_{0,m,n}\|_{L^2(\Omega^n)} \\ \implies \|\tilde{v}_m(t+h, \cdot) - \tilde{v}_m(t, \cdot)\|_{L^2(\Omega)} &\leq \frac{|h|}{t\sqrt{2}} \|u_{0,m}\|_{L^2(\Omega)}. \end{aligned} \quad (2.24)$$

Thus $\tilde{v}_m \in C([0, \infty); L^2(\Omega))$. By the contraction principle, $\tilde{v}_m = S^{\partial J_\Omega}(t)[u_{0,m}]$ is the unique generalized solution to (2.3). Finally, there exists an increasing sequence $\{u_{0,m}\} \subset L^2(\Omega)$ such that for any $\epsilon > 0$, and $\tau > 0$,

$$0 < \underline{u}_\Omega - S^{\partial J_\Omega}(t)[u_{0,m}] \leq \epsilon/2$$

on $[\tau, \infty) \times \Omega$. For any m , there exists n_m such that

$$0 < S^{\partial J_\Omega}(t)[u_{0,m}] - \tilde{v}_{m,n} \leq \epsilon/2$$

Therefore

$$0 < \underline{u}_\Omega - \underline{u}_{\Omega^n} \leq \epsilon,$$

on $[\tau, \infty) \times \Omega_n$. This implies (2.23). $\square$

We can also construct a minimal solution with conditional initial blow-up in the following way. Assuming that $\Omega = \cup \Omega^m$ where Ω^m are smooth bounded domains and $\overline{\Omega^m} \subset \Omega^{m+1}$. We denote by u_m the solution of

$$\begin{cases} \partial_t u_m - \Delta u_m + |u_m|^{q-1} u_m = 0 & \text{in } Q_\infty^{\Omega^m} \\ u_m = 0 & \text{in } \partial \Omega^m \times (0, \infty) \\ \lim_{t \rightarrow 0} u_m(x, t) = \infty & \text{locally uniformly on } \Omega^m. \end{cases} \quad (2.25)$$

Such a u_m is the increasing limit as $k \rightarrow \infty$ of the solutions $u_{m,k}$ of the same equation, with same boundary data and initial value equal to k . Since $\overline{\Omega^m} \subset \Omega^{m+1}$, $u_m < u_{m+1}$. We extend u_m by zero outside Ω^m and the limit of the sequence $\{u_m\}$, when $m \rightarrow \infty$ is a positive solution of (2.5) in Q_∞^Ω . We denote it by u_Ω . The next result is similar to Corollary 2.7, although the proof is much simpler.

Corollary 2.8 *Assume $\Omega^1 \subset \Omega^2 \subset \mathbb{R}^N$ are open domains, then $u_{\Omega^1} \leq u_{\Omega^2}$. Furthermore, if $\Omega = \cup \Omega^n$ where $\Omega^n \subset \Omega^{n+1}$, then*

$$\lim_{n \rightarrow \infty} u_{\Omega^n} = u_\Omega, \quad (2.26)$$

locally uniformly in Q_∞^Ω .

Proposition 2.9 *There holds $u_\Omega = \underline{u}_\Omega$.*

Proof. For any $m, k > 0$, $\tilde{u}_{m,k}$, the extension of $u_{m,k}$ by zero in $Q_\infty^{\Omega^{m,c}}$ is a subsolution, thus it is dominated by $\underline{u}_\Omega$. Letting successively $k \rightarrow \infty$ and $m \rightarrow \infty$ implies $u_\Omega \leq \underline{u}_\Omega$. In order to prove the reverse inequality, we consider an increasing sequence $\{u_\ell\} \subset \mathcal{I}(Q_\infty^\Omega)$ converging to $\underline{u}_\Omega$ locally uniformly in Q_∞^Ω . If Ω is bounded there exists a bounded sequence $\{u_{\ell,0,k}\}$ which converges to $u_\ell(\cdot, 0) = u_{\ell,0}$ in $L^2(\Omega)$ and $S^{\partial J_\Omega}(\cdot)[u_{\ell,0,k}] \rightarrow S^{\partial J_\Omega}(\cdot)[u_{\ell,0}]$ in $L^\infty(0, \infty; L^2(\Omega))$. Therefore

$$S^{\partial J_\Omega}(\cdot)[u_{\ell,0,k}] \leq u_\Omega \implies S^{\partial J_\Omega}(\cdot)[u_{\ell,0}] \leq u_\Omega \implies \underline{u}_\Omega \leq u_\Omega. \quad (2.27)$$

Next, if Ω is unbounded, $\Omega = \cup \Omega^n$, with $\Omega^n \subset \Omega^{n+1}$ are bounded, we have

$$\lim_{n \rightarrow \infty} \underline{u}_{\Omega^n} = \underline{u}_\Omega$$

and

$$\lim_{n \rightarrow \infty} u_{\Omega^n} = u_\Omega$$

by Corollary 2.7 and Corollary 2.8. Since $\underline{u}_{\Omega^n} = u_{\Omega^n}$ from the first part of the proof, the result follows. $\square$

Remark. By construction u_Ω is dominated by any positive solution of (2.12) which satisfies the initial blow-up condition locally uniformly in Ω . Therefore, $u_\Omega = \underline{u}_\Omega$ is the *minimal solution* with initial blow-up.

If Ω has the minimal regularity which allows the Dirichlet problem to be solved by any continuous function g given on $\partial\Omega \times [0, \infty)$, we can consider another construction of the maximal solution of (2.1) in Q_∞^Ω . The needed assumption on $\partial\Omega$ is known as the *parabolic Wiener criterion* [9] (abr. PWC).

Definition 2.10 *If $\partial\Omega$ is compact and satisfies PWC, we denote by $\mathcal{J}_{Q_\infty^\Omega}$ the set of $v \in C((0, \infty) \times \overline{\Omega}) \cap C^{2,1}(Q_\infty^\Omega)$ satisfying (2.1).*

Theorem 2.11 *Assume $q > 1$ and Ω satisfies PWC. Then $\mathcal{J}_{Q_\infty^\Omega}$ admits a maximal element $\overline{u}_\Omega$.*

Proof. Step 1- Construction. We shall directly assume that Ω is unbounded, the bounded case being a simple adaptation of our construction. We suppose $\Omega^c \subset B_{R_0}$, and for $n > R_0$ set $\Omega_n = \Omega \cap B_n$. The construction of u_n is standard: for $k \in \mathbb{N}_*$ we denote by $v_k^* = v_{n,k}^*$ the solution of (2.9). Lemma 2.5 is valid for v_k^* . Notice that uniqueness follows from the maximum principle. When $k \rightarrow \infty$ the sequence $\{v_k^*\}$ increases and converges to a solution u_n of (2.12) in Q_{Ω_n} . Because the exterior boundary of Ω_n is smooth, the standard equi-continuity of the sequence of solutions applies, thus $u_n(x, t) = 0$ for all (x, t) s.t. $|x| = n$ and $t > 0$. In order to see that $u_n(x, t) = 0$ for all (x, t) s.t. $x \in \partial\Omega$ and $t > 0$, we see that $u_n(x, t) \leq \phi_\tau(x, t)$ on $(\tau, \infty) \times \Omega_n$, where

$$\begin{cases} \partial_t \phi_\tau - \Delta \phi_\tau + \phi_\tau^q = 0 & \text{in } Q_\infty^\Omega \\ \phi_\tau(x, \tau) = \phi_q(\tau) & \text{in } \Omega \\ \phi_\tau(x, t) = 0 & \text{in } \partial\Omega \times [\tau, \infty) \end{cases} \quad (2.28)$$

Such a solution exists because of PWC assumption. Since $v_{n,k}^*$ is an increasing function of n (provided the solution is extended by 0 outside Ω_n) and k , there holds $\tilde{u}_n \leq u_{n+1}$ in Ω^{n+1} . If we set

$$\bar{u}_\Omega = \lim_{n \rightarrow \infty} \tilde{u}_n,$$

then $\bar{u}_\Omega \leq \phi_\tau$ for any $\tau > 0$. Clearly $\bar{u}_\Omega$ is a solution of (2.12) in Q_∞^Ω . This implies that $\bar{u}_\Omega$ is continuous up to $\partial\Omega \times (0, \infty)$, with zero boundary value. Thus it belongs to $\mathcal{J}_{Q_\infty^\Omega}$.

Step 2- Comparison. In order to compare $\bar{u}_\Omega$ to any other $u \in \mathcal{J}_{Q_\infty^\Omega}$, for $R > R_0$ we set $v_{R,\tau}(x, t) = \bar{u}_\Omega(x, t - \tau) + W_R(x)$, where W_R is the maximal solution of (2.20) in B_R . The function $(u - v_{R,\tau})_+$ is a subsolution of (2.12) in $\Omega \cap B_R \times (\tau, \infty)$. It vanishes in a neighborhood of $\partial(\Omega \cap B_R) \times (\tau, \infty)$ and of $\Omega \cap B_R \times \{\tau\}$. Thus it is identically zero. If we let $R \rightarrow \infty$ in the inequality $u \leq v_{R,\tau}$ and $\tau \rightarrow 0$, we derive $u \leq \bar{u}_\Omega$, which is the claim. $\square$

Proposition 2.12 *Under the assumptions of Theorem 2.11, $\bar{u}_\Omega \in \mathcal{I}(Q_\infty^\Omega)$ if Ω is bounded and $\bar{u}_\Omega \in \mathcal{I}(Q_\infty^{\Omega_{loc}})$ if Ω is not bounded.*

Proof. Case 1: Ω bounded. Let Ω^n be a sequence of smooth domains such that

$$\Omega^n \subset \bar{\Omega}^n \subset \Omega^{n+1} \subset \Omega$$

and $\cup_n \Omega^n = \Omega$. For $\tau > 0$, let $u_{n,\tau}$ be the solution of

$$\begin{cases} \partial_t u_{n,\tau} - \Delta u_{n,\tau} + u_{n,\tau}^q = 0 & \text{in } \Omega^n \times (\tau, \infty) \\ u_{n,\tau}(\cdot, \tau) = \bar{u}_\Omega(\cdot, \tau) & \text{in } \Omega^n \\ u_{n,\tau}(x, t) = 0 & \text{in } \partial\Omega^n \times [\tau, \infty) \end{cases} \quad (2.29)$$

Because $\bar{u}_\Omega(\cdot, \tau) \in C^2(\bar{\Omega}^n)$, $u_{n,\tau} \in C^{2,1}(\bar{\Omega}^n \times [\tau, \infty))$. By the maximum principle,

$$0 \leq \bar{u}_\Omega(\cdot, t) - u_{n,\tau}(\cdot, t) \leq \max\{\bar{u}_\Omega(x, s) : (x, s) \in \partial\Omega^n \times [\tau, t]\} \quad (2.30)$$

for any $t > \tau$. Because $\bar{u}_\Omega$ vanishes on $\partial\Omega \times [\tau, t]$, we derive

$$\lim_{n \rightarrow \infty} \tilde{u}_{n,\tau} = \bar{u}_\Omega \quad (2.31)$$

uniformly on $\bar{\Omega} \times [\tau, t]$ for any $t \geq \tau$, where $\tilde{u}_{n,\tau}$ is the extension of $u_{n,\tau}$ by zero outside Ω_n . Applying (2.15) and (2.19) with $\eta = 1$ to $\tilde{u}_{n,\tau}$ in Ω yields to

$$2^{-1} \int_\Omega \tilde{u}_{n,\tau}^2(\cdot, t) dx + \int_\tau^t \int_\Omega \left(|\nabla \tilde{u}_{n,\tau}|^2 + \tilde{u}_{n,\tau}^{q+1} \right) dx ds \leq C(t+1)\tau^{-2/(q-1)}. \quad (2.32)$$

and

$$2^{-1} \int_\tau^t \int_\Omega (s - \tau)(\partial_s \tilde{u}_{n,\tau})^2 dx ds + (t - \tau) \int_\Omega \left(\frac{|\nabla \tilde{u}_{n,\tau}|^2}{2} + \frac{\tilde{u}_{n,\tau}^{q+1}}{q+1} \right) (t, \cdot) dx \leq C(t - \tau)t\tau^{-2/(q-1)}. \quad (2.33)$$

Letting $n \rightarrow \infty$ and using (2.31) yields to

$$2^{-1} \int_{\Omega} \bar{u}_{\Omega}^2(., t) dx + \int_{\tau}^t \int_{\Omega} \left(|\nabla \bar{u}_{\Omega}|^2 + \bar{u}_{\Omega}^{q+1} \right) dx ds \leq C(t+1) \tau^{-2/(q-1)}. \quad (2.34)$$

and

$$2^{-1} \int_{\tau}^t \int_{\Omega} (s-\tau) (\partial_s \bar{u}_{\Omega})^2 dx ds + (t-\tau) \int_{\Omega} \left(\frac{|\nabla \bar{u}_{\Omega}|^2}{2} + \frac{\bar{u}_{\Omega}^{q+1}}{q+1} \right) (t, .) dx \leq C(t-\tau) t \tau^{-2/(q-1)}. \quad (2.35)$$

Since $L^2(\tau, t; H_0^1(\Omega))$ is a closed subspace of $L^2(\tau, t; H^1(\Omega))$, for any $0 < \tau < t$, $\bar{u}_{\Omega} \in L_{loc}^2(0, \infty; H_0^1(\Omega))$. Furthermore $\partial_s \bar{u}_{\Omega} \in L_{loc}^2(0, \infty; L^2(\Omega))$. Because $\bar{u}_{\Omega}$ satisfies (2.12), it implies $\bar{u}_{\Omega} \in \mathcal{I}(Q_{\infty}^{\Omega})$.

Case 2: Ω unbounded. We assume that $\Omega^c \subset B_{R_0}$. We consider a sequence of smooth unbounded domains $\{\Omega^n\} \subset \Omega$ ($n > 1$) such that $\sup\{\text{dist}(x, \Omega^c) : x \in \partial\Omega^n\} < 1/n$ as $n \rightarrow \infty$, thus $\cup_n \Omega^n = \Omega$. For $m > R_0$ we set $\Omega_m^n = \Omega^n \cap B_m$. Therefore $\Omega_m^n \subset \bar{\Omega}_m^n \subset \Omega_{m+1}^{n+1}$ and $\cup_{n,m} \Omega_m^n = \Omega$. For $\tau > 0$, let $u = u_{m,n,\tau}$ be the solution of

$$\begin{cases} \partial_t u - \Delta u + u^q = 0 & \text{in } \Omega_m^n \times (\tau, \infty) \\ u(., \tau) = \bar{u}_{\Omega}(., \tau) & \text{in } \Omega_m^n \\ u(x, t) = 0 & \text{in } \partial\Omega^n \times [\tau, \infty) \\ u(., \tau) = \bar{u}_{\Omega}(., \tau) & \text{in } \partial B_m \times (\tau, \infty). \end{cases} \quad (2.36)$$

By the maximum principle,

$$0 \leq \bar{u}_{\Omega}(., t) - u_{m,n,\tau}(., t) \leq \max\{\bar{u}_{\Omega}(x, s) : (x, s) \in \partial\Omega^n \times [\tau, t]\} \rightarrow 0, \quad (2.37)$$

as $n \rightarrow \infty$. Next we extend $u_{m,n,\tau}$ by zero in $\Omega \setminus \Omega_n$ and apply (2.15)-(2.19) with η as in Theorem 2.6 and $m > 2R$. We get, with $\Omega_R = \Omega \cap B_R$,

$$2^{-1} \int_{\Omega_R} u_{m,n,\tau}^2(., t) dx + \int_{\tau}^t \int_{\Omega_R} \left(2^{-1} |\nabla u_{m,n,\tau}|^2 + u_{m,n,\tau}^{q+1} \right) dx ds \leq CR^N(t+1) \tau^{-2/(q-1)}, \quad (2.38)$$

and

$$2^{-1} \int_{\tau}^t \int_{\Omega_R} (s-\tau) \left| \frac{du_{m,n,\tau}}{dt} \right|^2 dx ds + (t-\tau) \int_{\Omega_R} \left(\frac{|\nabla u_{m,n,\tau}|^2}{2} + \frac{u_{m,n,\tau}^{q+1}}{q+1} \right) dx \leq 8C(2R)^N(t-\tau) t \tau^{-2/(q-1)} \quad (2.39)$$

We let successively $m \rightarrow \infty$ and $n \rightarrow \infty$ and derive by Fatou's lemma and (2.37) that inequalities (2.38) and (2.39) still hold with $\bar{u}_{\Omega}$ instead of $u_{m,n,\tau}$. If we denote by $\tilde{H}_0^1(\Omega_R)$ the closure of the space of $C^\infty(\bar{\Omega}_R)$ functions which vanish in a neighborhood on $\partial\Omega$, then (2.38) is an estimate in $L^2(\tau, t; \tilde{H}_0^1(\Omega_R))$ which is a closed subspace of $L^2(\tau, t; H^1(\Omega_R))$. Therefore $\bar{u}_{\Omega} \in L_{loc}^2(0, \infty; \tilde{H}_0^1(\Omega_R))$. Using (2.39) and equation (2.12) we conclude that $\bar{u}_{\Omega} \in \mathcal{I}(Q_{\infty}^{\Omega_{loc}})$. $\square$

We end this section with a comparison result between $\underline{u}_{\Omega}$ and $\bar{u}_{\Omega}$.

Theorem 2.13 *Assume $q > 1$ and Ω satisfies PWC. Then $\underline{u}_\Omega = \overline{u}_\Omega$.*

Proof. By Proposition 2.9 and Theorem 2.11-Step 2, $\underline{u}_\Omega \leq \overline{u}_\Omega$. If Ω is bounded, we can compare $\underline{u}_\Omega(.,.)$ and $\overline{u}_\Omega(. + \tau, .)$ on $\Omega \times (0, \infty)$. Since $\underline{u}_\Omega$, the least upper bound of $\mathcal{I}(Q_\infty^\Omega)$ belongs to $\mathcal{I}(Q_\infty^\Omega)$, and $\overline{u}_\Omega(. + \tau, .) \in \mathcal{I}(Q_\infty^\Omega)$ we derive $\overline{u}_\Omega(. + \tau, .) \leq \underline{u}_\Omega(.,.)$, from which follows $\overline{u}_\Omega \leq \underline{u}_\Omega$. Next, if Ω is not bounded, we can proceed as in the proof of Theorem 2.6 by comparing $\underline{u}_\Omega(.,.) + W_R$ and $\overline{u}_\Omega(. + \tau, .)$ on $\Omega_R \times (0, \infty)$, where W_R is defined in (2.20). Because $(\overline{u}_\Omega(. + \tau, .) - \underline{u}_\Omega(.,.) - W_R(.))_+$ is a subsolution of (2.12) in $Q_\infty^{\Omega_R}$ which vanishes at $t = 0$ and near $\partial\Omega_R \times (0, \infty)$; it follows $\overline{u}_\Omega(. + \tau, .) \leq \underline{u}_\Omega(.,.) + W_R(.)$. Letting $R \rightarrow \infty$ and $\tau \rightarrow 0$ completes the proof. $\square$

3 Uniqueness of large solutions

Definition 3.1 *Let $q > 1$ and $\Omega \subset \mathbb{R}^N$ be any domain. A positive function $u \in C^{2,1}(Q_\infty^\Omega)$ of (2.12) is a large initial solution if it satisfies*

$$\lim_{t \rightarrow 0} u(x, t) = \infty \quad \forall x \in \Omega, \quad (3.40)$$

uniformly on any compact subset of Ω .

We start with the following lemma

Lemma 3.2 *Assume $u \in C^{2,1}(Q_\infty^\Omega)$ is a large solution of (2.12), then for any open subset G such that $\overline{G} \subset \Omega$, there holds*

$$\lim_{t \rightarrow 0} t^{1/(q-1)} u(x, t) = c_q := \left(\frac{1}{q-1} \right)^{1/(q-1)} \quad \text{uniformly in } G. \quad (3.41)$$

Proof. By compactness, it is sufficient to prove the result when $G = B_\rho$ and $\overline{B}_\rho \subset B_{\rho'} \subset \Omega$. Let $\tau > 0$; by comparison, $u(x, t) \geq u_{B_{\rho'}}(x, t + \tau)$ for any $(x, t) \in Q_\infty^\Omega$. Letting $\tau \rightarrow 0$ yields to $u \geq u_{B_{\rho'}}$. Next for $\tau > 0$,

$$\phi_q(t + \tau) \leq u_{B_{\rho'}}(x, t) + u_{B_{\rho'}^c}(x, t) + W_R(x) \quad \forall (x, t) \in Q_\infty^{\mathbb{R}^N}.$$

Similarly

$$\max\{u_{B_{\rho'}}(x, t + \tau), u_{B_{\rho'}^c}(x, t + \tau)\} \leq \phi_q(t) + W_R(x) \quad \forall (x, t) \in Q_\infty^{\mathbb{R}^N}.$$

Letting $R \rightarrow \infty$ and $\tau \rightarrow 0$,

$$\max\{u_{B_{\rho'}}, u_{B_{\rho'}^c}\} \leq \phi_q \leq u_{B_{\rho'}} + u_{B_{\rho'}^c} \quad \text{in } Q_\infty^{\mathbb{R}^N}.$$

For symmetry reasons, $x \mapsto u_{B_{\rho'}^c}(x, t)$ is radially increasing for any $t > 0$, thus, for any $\rho < \rho'$ and $T > 0$, there exists $C_{\rho, T} > 0$ such that

$$u_{B_{\rho'}^c}(x, t) \leq C_{\rho, T} \quad \forall (x, t) \in B_\rho \times [0, T].$$

Therefore

$$\lim_{t \rightarrow 0} t^{1/(q-1)} u_{B_{\rho'}}(x, t) = c_q \quad \text{uniformly on } B_\rho.$$

Because

$$u_{B_{\rho'}}(x, t) \leq u(x, t) \leq \phi_q(t) \quad \forall (x, t) \in Q_\infty^\Omega,$$

(3.41) follows. $\square$

As an immediate consequence of Lemma 3.2 and (2.23), we obtain

Proposition 3.3 *Assume $q > 1$ and $\partial\Omega$ is compact. Then u_Ω is a large solution.*

We start with the following uniqueness result

Proposition 3.4 *Assume $q > 1$, Ω satisfies PWC, $\partial\Omega$ is bounded, and either Ω or Ω^c is strictly starshaped with respect to some point. Then $\bar{u}_\Omega$ is the unique large solution belonging to $\mathcal{J}(Q_\infty^\Omega)$.*

Proof. Without loss of generality, we can suppose that either Ω or Ω^c is strictly starshaped with respect to 0. By Theorem 2.11, $\bar{u}_\Omega$ exists and, by (2.23) and Lemma 3.2, it is a large solution. Let $u \in \mathcal{J}(Q_\infty^\Omega)$ be another large solution. Clearly $u \leq \bar{u}_\Omega$. If Ω is starshaped, then for $k > 1$, the function $u_k(x, t) := k^{2/(q-1)} u(kx, k^2t)$ is a solution in Q_{Ω_k} , with $\Omega_k := k^{-1}\Omega$. Clearly it is a large solution and it belongs to $\mathcal{J}(Q_{\Omega_k})$. For $\tau \in (0, 1)$, set $u_{k,\tau}(x, t) = u_k(x, t - \tau)$. Because $\partial\Omega$ is compact,

$$\lim_{k \downarrow 1} d_H(\partial\Omega, \partial\Omega_k) = 0,$$

where d_H denotes the Hausdorff distance between compact sets. By assumption $\bar{u}_\Omega \in C([\tau, \infty) \times \bar{\Omega})$ vanishes on $[\tau, \infty) \times \partial\Omega$, thus, for any $\epsilon > 0$, there exists $k_0 > 1$ such that for any

$$k \in (1, k_0] \implies \sup\{\bar{u}_\Omega(x, t) : (x, t) \in [\tau, 1] \times \partial\Omega_k\} \leq \epsilon.$$

Since $u_{k,\tau} + \epsilon$ is a super solution in Q_{Ω_k} which dominates $\bar{u}_\Omega$ on $[\tau, 1] \times \partial\Omega_k$ and at $t = \tau$, it follows that $u_{k,\tau} + \epsilon \geq \bar{u}_\Omega$ in $(\tau, 1] \times \Omega_k$. Letting successively $k \rightarrow 1$, $\tau \rightarrow 0$ and using the fact that ϵ is arbitrary, yields to $u \geq \bar{u}_\Omega$ in $(0, 1] \times \Omega$ and thus in Q_∞^Ω . If Ω^c is starshaped, then the same construction holds provided we take $k < 1$ and use the fact that, for $R > 0$ large enough, $u_{k,\tau} + \epsilon + W_R$ is a super solution in $Q_{\Omega_k \cap B_R}$ which dominates $\bar{u}_\Omega$ on $[\tau, 1] \times \partial\Omega_k \cap B_R$ and at $t = \tau$. Letting successively $R \rightarrow \infty$, $k \rightarrow 1$, $\tau \rightarrow 0$ and $\epsilon \rightarrow 0$ yields to $u \geq \bar{u}_\Omega$. $\square$

As a consequence of Section 2, we have the more complete uniqueness theorem

Theorem 3.5 *Assume $q > 1$, $\Omega \subset \mathbb{R}^N$ is a domain with a bounded boundary $\partial\Omega$ satisfying PWC. Then for any $f \in C(\partial\Omega \times [0, \infty))$, $f \geq 0$, there exists a unique positive function $u = \bar{u}_{\Omega,f} \in C(\bar{\Omega} \times (0, \infty)) \cap C^{2,1}(Q_\infty^\Omega)$ satisfying*

$$\begin{cases} \partial_t u - \Delta u + |u|^{q-1}u = 0 & \text{in } Q_\infty^\Omega \\ u = f & \text{in } \partial\Omega \times (0, \infty) \\ \lim_{t \rightarrow 0} u(x, t) = \infty & \text{locally uniformly on } \Omega. \end{cases} \quad (3.42)$$

Proof. Step 1: Existence. It is a simple adaptation of the proof of Theorem 2.11. For $k, \tau > 0$, we denote by $u = u_{k,\tau,f}$ the solution of

$$\begin{cases} \partial_t u - \Delta u + |u|^{q-1}u = 0 & \text{in } \Omega \times (\tau, \infty) \\ u = f & \text{in } \partial\Omega \times (\tau, \infty) \\ u(x, \tau) = k & \text{on } \Omega. \end{cases} \quad (3.43)$$

Notice that $u_{k,\tau,f}$ is bounded from above by $\bar{u}_\Omega(\cdot, \cdot - \tau) + v_{f,\tau}$, where $v_{f,\tau} = v$ solves

$$\begin{cases} \partial_t v - \Delta v + |v|^{q-1}v = 0 & \text{in } \Omega \times (\tau, \infty) \\ v = f & \text{in } \partial\Omega \times (\tau, \infty) \\ v(x, \tau) = 0 & \text{on } \Omega. \end{cases} \quad (3.44)$$

If we let $k \rightarrow \infty$ we obtain a solution $u_{\infty,\tau,f}$ of the same problem except that the condition at $t = \tau$ becomes $\lim_{t \rightarrow \tau} u(x, t) = \infty$, locally uniformly for $x \in \Omega$. Clearly $u_{\infty,\tau,f}$ dominates in $\Omega \times (\tau, \infty)$ the restriction to this set of any $u \in C(\bar{\Omega} \times \infty) \cap C^{2,1}(Q_\infty^\Omega)$ solution of (3.42), in particular $\bar{u}_\Omega$. Therefore $u_{\infty,\tau,f} \geq u_{\infty,\tau',f}$ in $\Omega \times (\tau, \infty)$ for any $0 < \tau' < \tau$. When $\tau \rightarrow 0$, $u_{\infty,\tau,f}$ converges to a function $\bar{u}_f$ which satisfies the lateral boundary condition $\bar{u}_{\Omega,f} = f$. Therefore $\bar{u}_{\Omega,f}$ satisfies (3.42).

Step 2: Uniqueness. Assume that there exists another positive function $u := u_f \in C(\bar{\Omega} \times (0, \infty)) \cap C^{2,1}(Q_\infty^\Omega)$ solution of (3.42). Then $u_f < \bar{u}_{\Omega,f}$. For $\tau > 0$, consider the solution $v := v_\tau$ of

$$\begin{cases} \partial_t v - \Delta v + |v|^{q-1}v = 0 & \text{in } \Omega \times (\tau, \infty) \\ v = 0 & \text{in } \partial\Omega \times (\tau, \infty) \\ v(x, \tau) = u_f(x, \tau) & \text{on } \Omega. \end{cases} \quad (3.45)$$

Then $v_\tau \leq u_f$ in $\Omega \times (\tau, \infty)$. In the same way, we construct a solution $v : \tilde{v}_\tau$ of the same problem (3.45) except that the condition at $t = \tau$ is now $v(x, \tau) = \bar{u}_{\Omega,f}(x, \tau)$ for all $x \in \Omega$. Furthermore $v_\tau \leq \tilde{v}_\tau \leq \bar{u}_{\Omega,f}$. Next we adapt a method introduced in [6], [7] in a different context. We denote

$$Z_f = \bar{u}_{\Omega,f} - u_f \quad \text{and} \quad Z_{0,\tau} = \tilde{v}_\tau - v_\tau, \quad (3.46)$$

and, for $(r, s) \in \mathbb{R}_+^2$,

$$h(r, s) = \begin{cases} \frac{r^q - s^q}{r - s} & \text{if } r \neq s \\ 0 & \text{if } r = s. \end{cases}$$

Since $r \mapsto r^q$ is convex on $\mathbb{R}_+$, there holds

$$\begin{cases} r_0 \geq s_0, r_1 \geq s_1 \\ r_1 \geq r_0, s_1 \geq s_0 \end{cases} \implies h(r_1, s_1) \geq h(r_0, s_0).$$

This implies

$$h(u_{\Omega,f}, u_f) \geq h(\tilde{v}_\tau, v_\tau) \quad \text{in } \Omega \times [\tau, \infty). \quad (3.47)$$

Next we write

$$\begin{aligned} 0 &= \partial_t(Z_f - Z_{0,\tau}) - \Delta(Z_f - Z_{0,\tau}) + \bar{u}_{\Omega,f}^q - u_f^q - (\tilde{v}_\tau^q - v_\tau^q) \\ &= \partial_t(Z_f - Z_{0,\tau}) - \Delta(Z_f - Z_{0,\tau}) + h(\bar{u}_{\Omega,f}, u_f)Z_f - h(\tilde{v}_\tau, v_\tau)Z_{0,\tau}. \end{aligned} \quad (3.48)$$

Combining (3.47), (3.48) with the positivity of Z_f and $Z_{0,\tau}$, we derive

$$\partial_t(Z_f - Z_{0,\tau}) - \Delta(Z_f - Z_{0,\tau}) + h(\bar{u}_{\Omega,f}, u_f)(Z_f - Z_{0,\tau}) \leq 0, \quad (3.49)$$

in $\Omega \times (\tau, \infty)$. On $\partial\Omega \times [\tau, \infty)$ there holds $Z_f - Z_{0,\tau} = f - f = 0$. Furthermore, at $t = \tau$, $Z_f(x, \tau) - Z_{0,\tau}(x, \tau) = \bar{u}_{\Omega,f}(x, t) - u_f(x, \tau) - \bar{u}_{\Omega,f}(x, t) + u_f(x, \tau) = 0$. By the maximum principle, it follows $Z_f \leq Z_{0,\tau}$ in $\Omega \times [\tau, \infty)$. Since $\tau > \tau' > 0$ implies $v_\tau(x, \tau) = u_f(x, \tau) \geq v_{\tau'}(x, \tau)$ and $\tilde{v}_\tau(x, \tau) = \bar{u}_{\Omega,f}(x, \tau) \geq \tilde{v}_{\tau'}(x, \tau)$, the sequences $\{v_\tau\}$ and $\{\tilde{v}_\tau\}$ converge to some functions $\{v_0\}$ and $\tilde{v}_0$ which belong to $C(\bar{\Omega} \times (0, \infty)) \cap C^{2,1}(Q_\infty^\Omega)$ and satisfy (3.42) with $f = 0$ on $\partial\Omega \times (0, \infty)$. Furthermore

$$\bar{u}_{\Omega,f} - u_f \leq \tilde{v}_0 - v_0. \quad (3.50)$$

Since $\bar{u}_{\Omega,f} \geq \bar{u}_\Omega$, $\tilde{v}_0 \geq \bar{u}_\Omega$, which implies that $\tilde{v}_0 = \bar{u}_\Omega$ by the maximality of $\bar{u}_\Omega$. If Ω' is any smooth bounded open subset such that $\bar{\Omega}' \subset \Omega$ there holds by an easy approximation argument $v_0 \geq u_{\Omega'}$ in $\Omega' \times (0, \infty)$. Therefore $v_0 \geq u_\Omega = \underline{u}_\Omega = \bar{u}_\Omega$, by Proposition 2.9 and Theorem 2.13. Applying again Theorem 2.13 we derive that the right-hand side of (3.50) is zero, which yields to $\bar{u}_{\Omega,f} = u_f$ $\square$

References

- [1] W. Al Sayed and L. Véron, *On uniqueness of large solutions of nonlinear parabolic equations in nonsmooth domains*, Adv. Nonlinear Studies, to appear.
- [2] H. Brezis, *Propriétés régularisantes de certains semi-groupes non linéaires*, Isr. J. Math. **9**, 513-534 (1971).
- [3] H. Brezis, **Opérateurs maximaux monotones et semi-groupes de contractions dans des espaces de Hilbert**, North-Holland Mathematics Studies, **No. 5**. Notas de Matematica (50). North-Holland Publishing Co. (1973).
- [4] H. Brezis and A. Friedman, *Nonlinear parabolic equations involving measures as initial conditions*, J. Math. Pures Appl. **62**, 73-97 (1983).
- [5] J.B. Keller, *On solutions of $\Delta u = f(u)$* , Comm. Pure Appl. Math. **10**, 503-510 (1957).
- [6] M. Marcus and L. Véron, *The boundary trace of positive solutions of semilinear elliptic equations: the subcritical case*, Arch. Rat. Mech. Anal. **144**, 201-231 (1998).
- [7] M. Marcus and L. Véron, *The initial trace of positive solutions of semilinear parabolic equations*, Comm. Part. Diff. Equ. **24**, 1445-1499 (1999).
- [8] R. Osserman, *On the inequality $\Delta u \geq f(u)$* , Pacific J. Math. **7**, 1641-1647 (1957).
- [9] W. Ziemer, *Behavior at the boundary of solutions of quasilinear parabolic equations*, J. Differential Equations **35** 291-305 (1980).